

ARTICLE

Many Turán exponents via subdivisions

Tao Jiang^{1*} and Yu Qiu²

¹Department of Mathematics, Miami University, Oxford, OH 45056, USA and ²School of Mathematical Sciences, University of Science and Technology of China, Hefei, 230026, P.R. China

*Corresponding author. Email: jiangt@miamioh.edu

(Received 18 March 2021; revised 28 May 2022; accepted 23 June 2022; first published online 21 July 2022)

Abstract

Given a graph H and a positive integer n , the *Turán number* $\text{ex}(n, H)$ is the maximum number of edges in an n -vertex graph that does not contain H as a subgraph. A real number $r \in (1, 2)$ is called a *Turán exponent* if there exists a bipartite graph H such that $\text{ex}(n, H) = \Theta(n^r)$. A long-standing conjecture of Erdős and Simonovits states that $1 + \frac{p}{q}$ is a Turán exponent for all positive integers p and q with $q > p$.

In this paper, we show that $1 + \frac{p}{q}$ is a Turán exponent for all positive integers p and q with $q > p^2$. Our result also addresses a conjecture of Janzer [18].

Keywords: rational exponent conjecture; Turán numbers; subdivisions

2020 MSC Codes: 05C35

1. Introduction

1.1 Rational exponent conjecture

Given a family \mathcal{H} of graphs, the Turán number $\text{ex}(n, \mathcal{H})$ is the largest number of edges in an n -vertex graph that does not contain any member of \mathcal{H} as a subgraph. When \mathcal{H} consists of one single graph H , we write $\text{ex}(n, H)$ for $\text{ex}(n, \{H\})$.

Determining Turán numbers for various graphs is one of the central problems in extremal graph theory. The celebrated Erdős-Stone-Simonovits theorem states that for any non-bipartite graph H , $\text{ex}(n, H) = \left(1 - \frac{1}{\chi(H)-1}\right) \binom{n}{2} + o(n^2)$, where $\chi(H)$ is the chromatic number of H . For bipartite graphs H , it follows from the Kővári-Sós-Turán theorem that $\text{ex}(n, H) = O(n^{2-\alpha})$, where $\alpha = \alpha_H > 0$ is a constant. However, finding good estimates on $\text{ex}(n, H)$ for bipartite graphs H is difficult. Until recently, the order of magnitude of $\text{ex}(n, H)$ was known only for very few bipartite graphs H . Following [25], we say that a real number $r \in (1, 2)$ is *realizable* (by H) if there exists a bipartite graph H such that $\text{ex}(n, H) = \Theta(n^r)$. If r is realizable then we also call it a *Turán exponent*. A well-known conjecture of Erdős and Simonovits, known as the *rational exponent conjecture*, asserts that every rational number $r \in (1, 2)$ is a Turán exponent.

Conjecture 1.1 ([8]). *For all positive integers $q > p$, $1 + \frac{p}{q}$ is a Turán exponent.*

Until recently, the only rationals in $(1, 2)$ for which the conjecture was known to be true were rationals of the form $1 + \frac{1}{q}$ and $2 - \frac{1}{q}$ for positive integers $q \geq 2$, realized by so-called theta

[†]Research supported in part by NSF grant DMS-1855542.

[‡]Research supported in part by China Scholarship Council grant #201806340156.

graphs and complete bipartite graphs, respectively. In a recent breakthrough work, Bukh and Conlon [3] showed that for any rational number $r \in (1, 2)$, there exists a finite family \mathcal{H}_r of graphs such that $\text{ex}(n, \mathcal{H}_r) = \Theta(n^r)$. Bukh and Conlon's work has, to a large extent, rejuvenated people's interest on Conjecture 1.1. In the last year or so, several new infinite sequences of new Turán exponents have been obtained by various groups. First, Jiang, Ma, and Yepremyan [21] showed that $2 - \frac{2}{2m+1}$ is realizable by generalized cubes and that $\frac{7}{5}$ is realizable by the so-called 3-comb-pasting graph. A few months later, Kang, Kim, and Liu [25] showed that for all positive integers $p < q$, where $q \equiv \pm 1 \pmod{p}$, $2 - \frac{p}{q}$ is realizable. More specifically, rationals of the form $2 - \frac{t}{st-1}$, where $s, t \geq 2$, are realized by the so-called *blowups* of certain height 2 trees. (We will define blowups precisely in subsection 1.2.) Rationals of the form $2 - \frac{t}{st+1}$ are realized by graphs obtained from theta graphs via some iterative operations. More recently, some new sequences of Turán exponents were obtained along the study of Turán numbers of subdivisions. For any integers $s, t \geq 1, k \geq 2$, let $K_{s,t}^k$ denote the graph obtained from the complete bipartite graph $K_{s,t}$ by subdividing each edge $k - 1$ times. Let $L_{s,t}(k)$ be obtained from $K_{s,t}^k$ by adding an extra vertex joined to all vertices in the part of $K_{s,t}$ of size t . Confirming a conjecture of Kang, Kim, and Liu [25], Conlon, Janzer, and Lee [7] showed that there exists t_0 such that for all integers $s, k \geq 1, t \geq t_0$, $\text{ex}(n, L_{s,t}(k)) = \Theta\left(n^{1+\frac{s}{sk+1}}\right)$, and thus establishing $1 + \frac{s}{sk+1}$ as Turán exponents. Subsequently, in verifying a conjecture of Conlon, Janzer, and Lee [7], Janzer [18] proved that there exists a t_0 such that for all integers $s, k \geq 2, t \geq t_0$, $\text{ex}(n, K_{s,t}^k) = \Theta\left(n^{1+\frac{s-1}{sk}}\right)$, thus establishing $1 + \frac{s-1}{sk}$ as Turán exponents. Earlier, Conlon, Janzer, Lee [7] had proven the conjecture for $k = 2$, while Jiang and Qiu [22] proved the conjecture for $k = 3, 4$.

1.2 Our results

In this paper, we build on the recent work on subdivisions to establish the following large three-parameter family of Turán exponents, which includes all the ones obtained by Conlon, Janzer, and Lee [7] and by Janzer [18].

Theorem 1.2. *For any positive integers p, k, b with $k \geq b$, $1 + \frac{p}{kp+b}$ is a Turán exponent.*

As an immediate corollary, we get the following easily stated result.

Theorem 1.3. *For any positive integers p and q with $q > p^2$, $1 + \frac{p}{q}$ is a Turán exponent.*

Using a reduction lemma of Kang, Kim, and Liu [25], Theorem 1.2 also yields

Corollary 1.4. *For any integers $b, p, s \geq 1$ and $k \geq 0$, if $k \geq b - 1$, then $2 - \frac{kp+b}{s(kp+b)+p}$ is a Turán exponent.*

Corollary 1.4 implies the following.

Corollary 1.5. *For any positive integers p, q with $q > p$, if $(q \bmod p) \leq \sqrt{p}$, then $2 - \frac{p}{q}$ is a Turán exponent.*

Theorem 1.2 follows from a theorem (Theorem 1.15) that we prove on the Turán number of subdivisions of $K_{s,t}$ where different edges of $K_{s,t}$ may be subdivided different number of times. The theorem is interesting on its own and partially answers a conjecture of Janzer (Conjecture 1.13), which we will describe in the next subsection.

1.3 The Bukh-Conlon Conjecture and Janzer's conjecture

At the core of the work of Bukh and Conlon [3] is the study of so-called blowups of balanced rooted trees, defined as follows (also see [3]).

Definition 1.6. A rooted tree (T, R) consists of a tree T together with an independent set $R \subseteq V(T)$, which we refer to as the roots. When the choice of R is clear, we will simply write T for (T, R) .

Definition 1.7. Given a rooted tree (T, R) and a non-empty subset $S \subseteq V(T) \setminus R$, let $\rho_T(S) = \frac{e(S)}{|S|}$, where $e(S)$ is the number of edges in T that have at least one end in S . Let $\rho_T = \rho_T(V(T) \setminus R)$ and call it the density of T . We say (T, R) is balanced if $\rho_T(S) \geq \rho(T)$ for any non-empty subset $S \subseteq V(T) \setminus R$.

Definition 1.8. The t -blowup of a rooted tree (T, R) , denoted by $t * T_R$, is the union of t labelled copies of T which agree on R but are pairwise vertex disjoint outside R . If the choice of R is clear, then we write $t * T$ for $t * T_R$.

The key result of Bukh and Conlon [3] is the following lower bound theorem, established using an innovative random algebraic approach. Interested readers can find the full statement in [3].

Theorem 1.9 ([3]). *Suppose that (T, R) is a balanced rooted tree with density ρ . Then there exists an integer $t_0 \geq 2$ such that for all integers $t \geq t_0$ we have $\text{ex}(n, t * T_R) = \Omega\left(n^{2-\frac{1}{\rho}}\right)$.*

Bukh and Conlon further made the following conjecture on a matching upper bound.

Conjecture 1.10 ([3]). *Suppose that (T, R) is a balanced rooted tree with density ρ . Then for all positive integers t we have $\text{ex}(n, t * T_R) = O\left(n^{2-\frac{1}{\rho}}\right)$.*

Besides being interesting on its own, a significance of Conjecture 1.10 is that it implies the rational exponent conjecture. Indeed, for each rational $r \in (1, 2)$, Bukh and Conlon were able to construct a balanced rooted tree (T, R) with density $\rho = \frac{1}{2-r}$. Hence Theorem 1.9 and Conjecture 1.10 together would give $\text{ex}(n, t * T_R) = \Theta(n^t)$ for some sufficiently large positive integer t . A careful reader will note that Bukh and Conlon’s conjecture is in fact much stronger than the rational exponent conjecture. Indeed, to prove the rational exponent conjecture, it suffices to find, for each $r \in (1, 2)$, a balanced rooted tree (T, R) with density $\rho = \frac{1}{2-r}$ for which the Bukh-Conlon conjecture holds. This suggests that one way to make further progress on the rational exponent conjecture is to find suitable balanced rooted trees to explore Conjecture 1.10 with. One family of trees whose exploration has brought some success are the so-called spiders.

Definition 1.11. Let $s \geq 2$ be an integer. An s -legged spider S with centre u is a tree consisting of s paths (called the legs of S) that share one common end u but are vertex disjoint outside u . Moreover, we say S has length vector (j_1, \dots, j_s) and leaf vector (x_1, \dots, x_s) if for every $1 \leq i \leq s$, its i -th leg has length j_i and has ends u and x_i .

For spiders with roots being all of its leaves, checking balancedness is simple.

Proposition 1.12. *Let s, k be integers where $s \geq 2, k \geq 1$. Let S be an s -legged spider and R the set of its leaves. Suppose the longest leg of S has length k . Then (S, R) is a balanced rooted tree if and only if $e(S) \geq (s - 1)k$.*

When S is an s -legged spider with length vector (k, \dots, k) and R is the set of its leaves, $t * S_R$ is the subdivision $K_{s,t}^k$ of $K_{s,t}$, considered by Janzer [18]. When S is an $(s + 1)$ -legged spider with length vector $(1, k, \dots, k)$ and R is the set of its leaves, $t * S_R$ is the graph $L_{s,t}(k)$, considered by Conlon, Janzer, and Lee [7]. Motivated by the earlier mentioned results on $\text{ex}(n, L_{s,t}(k))$ and $\text{ex}(n, K_{s,t}^k)$, Janzer [18] made the following conjecture.

Conjecture 1.13 ([18]). *Let $s \geq 2, k, b, t \geq 1$ be integers. Let S be an s -legged spider where the longest leg has length k . Suppose that $e(S) = (s - 1)k + b$, where $0 \leq b \leq k$. Then $\text{ex}(n, t * S) = O\left(n^{1+\frac{s-1}{(s-1)k+b}}\right)$.*

Even though Janzer’s conjecture is a special case of the Bukh-Conlon conjecture, it is also interesting on its own due to its connection to the study of subdivisions. Let S be as specified in Conjecture 1.13. It follows from Theorem 1.9 that there exists a t_0 such that for all $t \geq t_0$, $\text{ex}(n, t * S) = \Omega\left(n^{1 + \frac{s-1}{(s-1)k+b}}\right)$. Hence, if Conjecture 1.13 is true, it will establish all rationals of the form $1 + \frac{p}{pk+b}$ as Turán exponents, where p, k are positive integers and b is an integer with $0 \leq b \leq k$. Here, we settle an important case of Conjecture 1.13 that allows us to obtain all the Turán exponents that Conjecture 1.13 would give.

Definition 1.14. For positive integers k, b and s , let $S_{b,k}^s$ denote the s -legged spider with length vector (b, k, \dots, k) .

Using this notation, we have $K_{s,t}^k = t * S_{k,k}^s$ and $L_{s,t}(k) = t * S_{1,k}^{s+1}$. In this paper, we will prove the following common generalization of the result of Conlon, Janzer, and Lee on $\text{ex}(n, L_{s,t}(k))$ and the result of Janzer on $\text{ex}\left(n, K_{s,t}^k\right)$, from which our main theorem, Theorem 1.2, follows.

Theorem 1.15. For any $s, t \geq 2$ and $k \geq b \geq 1$, $\text{ex}\left(n, t * S_{b,k}^s\right) = O\left(n^{1 + \frac{s-1}{(s-1)k+b}}\right)$.

As in [6, 7, 22, 18], we will use the following variant of the regularization lemma of Erdős and Simonovits [11], as given in [24]. Given a positive constant K , a graph G is K -almost-regular if $\Delta(G) \leq K\delta(G)$.

Lemma 1.16 ([24]). Let $0 < \epsilon < 1$ and $c \geq 1$. There exists $n_0 = n_0(\epsilon) > 0$ such that the following holds for all $n \geq n_0$. If G is a graph on n vertices with $e(G) \geq cn^{1+\epsilon}$, then G contains a K -almost-regular subgraph G' on $m \geq n^{\frac{\epsilon-2}{2+2\epsilon}}$ vertices such that $e(G') \geq \frac{2c}{5}m^{1+\epsilon}$ and $K = \left\lceil 20 \cdot 2^{\frac{1}{\epsilon^2}+1} \right\rceil$.

By Lemma 1.16, in order to prove Theorem 1.15, it suffices to prove the following.

Theorem 1.17. Let $s, t \geq 2$ and $k \geq b \geq 1$. Let $K = K(s, b, k)$ be obtained by Lemma 1.16 with $\epsilon := \frac{s-1}{(s-1)k+b}$. There exist positive constants n_0 and C depending only on s, t, b, k such that for all integers $n \geq n_0$ if G is an n -vertex $t * S_{b,k}^s$ -free K -almost-regular graph then $\delta(G) < Cn^{\frac{s-1}{(s-1)k+b}}$.

The rest of the paper is organized as follows. In Section 2, we introduce some notation and preliminary lemmas. In Section 3, we prove Theorem 1.17, from which Theorems 1.15 and 1.2 follow. In Section 4, we give a sketch of proofs of Corollaries 1.4 and 1.5 and some concluding remarks.

2. Notation and preliminaries

Given a positive integer m , let $[m] = \{1, \dots, m\}$. Given a graph G and a vertex w , for each $i \geq 1$ let $\Gamma_i(w)$ be the set of vertices z such that there exists a path in G of length i with ends w and z . When $i=1$, we often write $N_G(w)$ for $\Gamma_1(w)$. Let $e(G)$ be the number of edges in G . We use standard asymptotic notations, that is, given two positive functions $f(n)$ and $g(n)$, by $f = o_n(g), f = \omega_n(g), f = \Omega_n(g), f = O_n(g), f = \Theta_n(g)$, we respectively mean $\lim_{n \rightarrow \infty} f/g = 0, \liminf_{n \rightarrow \infty} f/g = \infty, \liminf_{n \rightarrow \infty} f/g > 0, \limsup_{n \rightarrow \infty} f/g < \infty, 0 < \liminf_{n \rightarrow \infty} f/g \leq \limsup_{n \rightarrow \infty} f/g < \infty$. Whenever the context is clear, we drop the subscript n . If G is a graph and S is a set of vertices in it, then we define

$$N_G^*(S) = \bigcap_{x \in S} N_G(x),$$

and call it the *common neighbourhood* of S in G .

For the rest of the paper, we fix integers $s, t \geq 2$ and $k \geq b \geq 1$, and let $K = K(s, b, k)$ be obtained by Lemma 1.16 with $\epsilon := \frac{s-1}{(s-1)k+b}$.

Below are some key concepts introduced in [7], which we adapt for our setting.

Definition 2.1. Let L be a positive integer, we define $f(1, L) = L$ and for $j \geq 2$,

$$f(j, L) := 10j^4 [2K^j L \cdot f(j-1, L)^2]^{s+3}.$$

We will need the following property of the function in various places of the paper.

Proposition 2.2. For every integer $j \geq 2$, $\frac{f(j, L)}{j^2 f(j-1, L)^2} \geq \max\{2L^2, f(j-1, L)\}$ holds.

The next two definitions are crucial to our overall arguments.

Definition 2.3. We recursively define j -admissible, j -light paths, and j -heavy paths in a graph G . Any edge is both 1-admissible and 1-light. For $j \geq 2$, a path P is j -admissible if it has length j and for each $1 \leq \ell < j$ every subpath of length ℓ in P is ℓ -light.

Among j -admissible paths P with ends x and y , we further say that P is j -light if the number of j -admissible paths with ends x and y in G is less than $f(j, L)$ and that P is j -heavy otherwise.

Since the length of a path P is fixed, we often drop the prefix j and ℓ in the definitions above. Note that j -admissible and j -light paths are defined for all $j \geq 1$ while j -heavy paths are defined only for $j \geq 2$. In [22], the concepts of admissible, light, and heavy paths were extended for spiders. Here, we adapt the definitions from [22] further.

Definition 2.4. We recursively define s -legged admissible, light, and heavy spiders in a graph G . Any spider of height 1 is both admissible and light. Let S be an s -legged spider with leaf vector (x_1, \dots, x_s) and length vector $(j_1, \dots, j_s) \neq (1, 1, \dots, 1)$. We say that S is admissible if every leg of it is a light path as defined in Definition 2.3 and every s -legged proper sub-spider of S is light. Suppose S is admissible. Then we further say that it is light if the number of admissible spiders in G with leaf vector (x_1, \dots, x_s) and length vector (j_1, \dots, j_s) is less than $f(j, L)$ where $j = j_1 + \dots + j_s$. If S is admissible but not light, then we say that it is heavy.

At this point, let us say a few words about the function $f(j, L)$ given in Definition 2.1, as this function plays an important role in our arguments. In application we always assume that the parameter L is sufficiently larger than s, t, k and K and roughly speaking $f(j, L)$ is chosen so that $f(j, L) \gg f(j-1, L)$, that is, $\frac{f(j, L)}{f(j-1, L)} \rightarrow \infty$ as $L \rightarrow \infty$.

Next, we give several lemmas. Lemma 2.6 is similar to one used in [7]. Lemma 2.7 has its analogous counterparts in [22] and [18]. However, since our terminologies and choices of constants are slightly different, we include full proofs for completeness.

Lemma 2.5. Let G be a K -almost-regular graph. Let $1 \leq i \leq j$ be integers. Let x, w, y be vertices in G . Then the number of j -admissible paths in G that have x, w, y as the first, $(i+1)$ -th and last vertices, respectively is at most $f(i, L) \cdot f(j-i, L)$. Furthermore, if $i = 1$ or $j-1$, then there are most $f(j-1, L)$ such paths.

Proof. Let \mathcal{P} be the family of j -admissible paths in G that have x, w, y as the first, $(i+1)$ -th, and last vertices, respectively. Let $P \in \mathcal{P}$, by definition, each proper subpath of P is light. So P is the union of an i -light path from x to w and a $(j-i)$ -light path from w to y . By definition of light paths there are at most $f(i, L)$ i -light paths in G with ends x, w and at most $f(j-i, L)$ $(j-i)$ -light paths with ends w and y . So $|\mathcal{P}| \leq f(i, L) \cdot f(j-i, L)$.

If $i = 1$ then every $P \in \mathcal{P}$ is the union of the edge xw and a $(j-1)$ -light path with ends w and y . So $|\mathcal{P}| \leq f(j-1, L)$. The case $i = j-1$ is similar. \square

Lemma 2.6. Let x, y be two vertices and \mathcal{C} be a family of j -admissible paths between x and y . Then there are at least $|\mathcal{C}|/j^2 \cdot f(j-1, L)^2$ members of \mathcal{C} that are pairwise vertex disjoint outside $\{x, y\}$.

Proof. Let $\mathcal{C}' = \{Q_1, \dots, Q_r\} \subseteq \mathcal{C}$ be a maximal subfamily of \mathcal{C} that are pairwise vertex disjoint outside $\{x, y\}$. Let $W = \bigcup_{i=1}^r V(Q_i) \setminus \{x, y\}$. Then $|W| = (j - 1)r$. By maximality, every member of \mathcal{C} must contain a vertex $v \in W$ as an internal vertex. For each $v \in W$ and each $1 \leq i \leq j - 1$ let $\mathcal{C}_{v,i}$ be the subfamily of members of \mathcal{C} that contain v as its $(i + 1)$ -th vertex (when the member is viewed from x to y). Then $\mathcal{C} = \bigcup_{v,i} \mathcal{C}_{v,i}$. By Lemma 2.5, for any fixed v, i , we have $|\mathcal{C}_{v,i}| \leq f(i, L) \cdot f(j - i, L) \leq f(j - 1, L)^2$. Hence

$$|\mathcal{C}| = \left| \bigcup_{v,i} \mathcal{C}_{v,i} \right| \leq \sum_{v \in W} \sum_{i=1}^{j-1} f(j - 1, L)^2 < rj^2 f(j - 1, L)^2.$$

Solving the inequality for r , we get the desired claim. □

For two spiders with the same leaf vector and length vector, we say they are *internally disjoint* if they are vertex disjoint outside their leaves.

Lemma 2.7. *Let \mathcal{S} be a family of admissible spiders with leaf vector (x_1, \dots, x_s) and length vector (j_1, \dots, j_s) . Then there are at least $|\mathcal{S}| / [j^2 \cdot f(j - 1, L)^2]$ members of \mathcal{S} that are pairwise vertex disjoint outside $\{x_1, \dots, x_s\}$, where $j = j_1 + \dots + j_s$.*

Proof. Let $\mathcal{S}' = \{S_1, \dots, S_r\} \subseteq \mathcal{S}$ be a maximal subfamily of members of \mathcal{S} that are pairwise vertex disjoint outside $\{x_1, \dots, x_s\}$. Let $W = \bigcup_{i=1}^r V(S_i) \setminus \{x_1, \dots, x_s\}$. Then $|W| = (j - s + 1)r$. By maximality of \mathcal{S}' , every member of \mathcal{S} must contain some $v \in W$ as a non-leaf vertex. For each $v \in W$ let \mathcal{D}_v denote the subfamily of members of \mathcal{S} that contain v as the centre. For each $v \in W$, $i \in [s]$, and $1 \leq \ell < j_i$, let $\mathcal{S}_{v,i,\ell}$ denote the subfamily of members of \mathcal{S} in which v is on the i -th leg and the distance from v to x_i is ℓ . Then $\mathcal{S} = (\bigcup_{v \in W} \mathcal{D}_v) \cup (\bigcup_{v \in W, i \in [s], 1 \leq \ell < j_i} \mathcal{S}_{v,i,\ell})$.

Let $S \in \mathcal{D}_v$. Then by definition, for each $i \in [s]$, the i -th leg of S is a j_i -light path between v and x_i . Hence, by the definition of light paths,

$$|\mathcal{D}_v| \leq \prod_{i=1}^s f(j_i, L) \leq f(j - 1, L)^2,$$

where the last inequality holds because by Definition 2.1 we have that $\prod_{i=1}^s f(j_i, L) \leq f(j_1 + j_2 - 1, L)^2 \prod_{i=3}^s f(j_i, L) \leq f(j_1 + j_2, L) \prod_{i=3}^s f(j_i, L) \leq \dots \leq f(j_1 + \dots + j_{s-1}, L) f(j_s, L) \leq f(j - 1, L)^2$.

Next, fix $v \in W$, $i \in [s]$, and $1 \leq \ell < j_i$. Let $S \in \mathcal{S}_{v,i,\ell}$. Since S is admissible, the v, x_i -path in S is ℓ -light while the rest of S is an s -legged proper sub-spider, which by definition, is light. This implies that

$$|\mathcal{S}_{v,i,\ell}| \leq f(\ell, L) f(j - \ell, L) \leq f(j - 1, L)^2.$$

Putting everything together, we obtain

$$|\mathcal{S}| \leq |W| f(j - 1, L)^2 + \sum_{v \in W} \sum_{i=1}^s \sum_{\ell=1}^{j_i-1} f(j - 1, L)^2,$$

which implies that $|\mathcal{S}| \leq rj^2 f(j - 1, L)^2$, from which the claim follows. □

The following lemma is proved in [22]. A spider has *height* ℓ if all of its legs have length ℓ .

Lemma 2.8 ([22] Lemma 3.6). *Let G be a K -almost-regular graph with minimum degree δ . Let x be a vertex. Let \mathcal{C} be a family of paths of length h with one end x and another end in a set S . For each $i \in [h]$ there exists a vertex x_i and a spider of height i with centre x_i and leaves in S which has at least $|\mathcal{C}| / [h(K\delta)^{h-1}]$ legs. Furthermore, $x_i = x$ if and only if $i = h$.*

We also need a standard averaging lemma as below.

Lemma 2.9. *Let $0 < c < 1$ be a real and m be a positive integer. Let G be a bipartite graph with a bipartition (X, Y) . Suppose that $e(G) \geq c|X||Y|$ and that $c|X| \geq 2m$. Then there exists an m -set S in X such that $|N_G^*(S)| \geq (c/2)^m|Y|$.*

Proof. By our assumption, the average degree of vertices in Y is at least $c|X|$. Let \mathcal{F} be the family of $K_{1,m}$'s with centre in Y . Then $|\mathcal{F}| = \sum_{y \in Y} \binom{d_G(y)}{m} \geq |Y| \binom{c|X|}{m}$, where the last inequality uses the convexity of the function $\binom{x}{m}$. Hence, by averaging there exists an m -set S in X such that the number of members of \mathcal{F} that have S as the leaf set is at least

$$|Y| \frac{\binom{c|X|}{m}}{\binom{|X|}{m}} \geq |Y| \left(\frac{c|X| - m}{|X| - m} \right)^m > (c/2)^m|Y|,$$

where the last inequality uses the condition $c|X| \geq 2m$. □

Finally, we need a standard cleaning lemma.

Lemma 2.10. *If B is a bipartite graph with parts X and Y , then it has subgraph B' such that $e(B') \geq \frac{e(B)}{2}$ and $\forall x \in X \cap V(B'), d_{B'}(x) \geq \frac{e(B)}{4|X|}$ and $\forall y \in Y \cap V(B'), d_{B'}(y) \geq \frac{e(B)}{4|Y|}$.*

Proof. Whenever there is a vertex in X whose degree becomes less than $\frac{e(B)}{4|X|}$ or a vertex in Y whose degree becomes less than $\frac{e(B)}{4|Y|}$, we delete it. Let B' denote the final subgraph of B . As the number of edges deleted is at most $|X| \cdot \frac{e(B)}{4|X|} + |Y| \cdot \frac{e(B)}{4|Y|} = \frac{e(B)}{2}$, $e(B') \geq \frac{e(B)}{2}$. By definition, B' satisfies our requirements. □

3. Proof of Theorem 1.17

3.1 Overall structure of the proof

Our overall strategy has roots in the work of Conlon and Lee [6] and the work of Conlon, Janzer, and Lee [7], particularly [7]. Some of the strategies used there were later augmented (through the concepts of admissible, light, and heavy spiders) in the work of Jiang and Qiu [22] and the work of Janzer [18]. In particular, Janzer [18] introduced a creative way to extending spiders, an idea that we will develop further. Overall, our proof combines ideas from [7], [22], [18] and some new ideas.

Let G be a K -almost-regular $t * S_{b,k}^s$ -free graph on n vertices, where n is sufficiently large. To prove the theorem, it suffices to show that there exists a constant C depending on s, b, k such that if $\delta(G) \geq Cn^{\frac{s-1}{(s-1)k+b}}$, then G must contain a copy of $t * S_{b,k}^s$, which would contradict G being $t * S_{b,k}^s$ -free and complete the proof. The general strategy is to show that (1) G contains many copies of $S_{b,k}^s$ and (2) most of these copies of $S_{b,k}^s$ are light. Then by averaging, there exist some vector (x_1, \dots, x_s) of s vertices which is the leaf vector of a large number of light copies of $S_{b,k}^s$. This will imply that all these spiders are heavy, giving us contradiction. More specifically, the proof of Theorem 1.17 follows readily after we establish the following two crucial lemmas.

Lemma 3.1. *Let G be a $t * S_{b,k}^s$ -free K -almost-regular graph on n vertices with minimum degree $\delta = \omega(1)$. Then provided that L is sufficiently large compared to s, t, k, K , for any $2 \leq j \leq k$, the number of j -heavy paths in G is at most $\frac{(j+1)^{j+1}}{L} n \delta^j$.*

Lemma 3.2. *Let G be a $t * S_{b,k}^s$ -free K -almost-regular graph on n vertices with minimum degree $\delta = \omega(1)$. Let $1 \leq j_1 \leq b$ and $1 \leq j_2, \dots, j_s \leq k$ be integers. Then provided that L is sufficiently large*

compared to s, t, k, K , the number of heavy spiders with length vector (j_1, \dots, j_s) is at most $\frac{27K^{j-2}}{L} n\delta^j$ where $j = j_1 + \dots + j_s$.

We now show how Theorem 1.17 follows from Lemma 3.1 and Lemma 3.2.

Proof of Theorem 1.17. Let L be a sufficiently large constant compared to s, t, k, K . Let G be a K -almost-regular $t * S_{b,k}^s$ -free graph on n vertices with minimum degree δ . Let $h = e(S_{b,k}^s) = (s - 1)k + b$. Suppose to the contrary that $\delta \geq Cn^{\frac{s-1}{(s-1)k+b}}$, where $C := 2f(h, L)(h + 1)!$. Let \mathcal{S} be the family of spiders in G with length vector (b, k, \dots, k) . By a greedy process, it is easy to see that

$$|\mathcal{S}| \geq \frac{1 - o(1)}{(h + 1)!} \cdot n\delta^h.$$

Let \mathcal{S}_1 be the family of spiders in \mathcal{S} that contain some heavy path of length $2 \leq j \leq k$. As the maximum degree of G is at most $K\delta$, by Lemma 3.1, we have

$$|\mathcal{S}_1| \leq \sum_{j=2}^k \binom{h}{j} \frac{(j + 1)^{j+1}}{L} n\delta^j (K\delta)^{h-j} \leq \frac{(k + 1)^{k+2} K^{sk} h!}{L} n\delta^h,$$

where the factor $\binom{h}{j}$ upper bounds the number of positions of a j -heavy path in $S_{b,k}^s$. Let \mathcal{S}_2 be the family of spiders in \mathcal{S} that contain some s -legged heavy sub-spider. As the maximum degree of G is at most $K\delta$, by Lemma 3.2, we have

$$|\mathcal{S}_2| \leq \sum_{\substack{1 \leq j_1 \leq b \\ 1 \leq j_2, \dots, j_s \leq k}} \frac{27K^{j_1 + \dots + j_s - 2}}{L} n\delta^{j_1 + \dots + j_s} \cdot (K\delta)^{h - (j_1 + \dots + j_s)} \leq \frac{27K^{sk} k^s}{L} n\delta^h.$$

Let $\mathcal{S}' = \mathcal{S} - (\mathcal{S}_1 \cup \mathcal{S}_2)$. Then it follows that

$$|\mathcal{S}'| \geq |\mathcal{S}| - (|\mathcal{S}_1| + |\mathcal{S}_2|) \geq \frac{1 - o(1)}{(h + 1)!} \cdot n\delta^h - \frac{K^{sk}(27k^s + (k + 1)^{k+2}h!)}{L} n\delta^h \geq \frac{n\delta^h}{2(h + 1)!},$$

where the last inequality holds since L is sufficiently large. As $\delta \geq Cn^{\frac{s-1}{(s-1)k+b}}$, and $C = 2(h + 1)f(h, L)$, it follows that $|\mathcal{S}'| \geq f(h, L)n^s$. By averaging, there exists an s -tuple (x_1, \dots, x_s) of distinct vertices, such that the subfamily \mathcal{S}'' which consists of all spiders in \mathcal{S}' with leaf vector (x_1, \dots, x_s) has size $|\mathcal{S}''| \geq f(h, L)$. For any $S \in \mathcal{S}''$, since S contains no heavy path of length at most k , every leg of S is light. Since S does not contain any s -legged heavy sub-spider, S is light. So \mathcal{S}'' is a family of at least $f(h, L)$ light spiders with leaf vector (x_1, \dots, x_s) and length vector (b, k, \dots, k) . This contradicts the definition of the light spider with length vector (b, k, \dots, k) . \square

Thus, to complete our proof of Theorem 1.17, it remains to prove Lemma 3.1 and Lemma 3.2. Lemma 3.2 was proved by Janzer for the case $b = k$ in details in [18]. An outline of the general case was given in Lemma 4.3 in the same paper. Since our Lemma 3.2 is essentially the same as Lemma 4.3 of [18] apart from some technical details, we omit its proof. Readers who are interested in the details of the proof of Lemma 3.2 are referred to the arxiv version of the manuscript [23]. As the author of [18] pointed out the main obstacle to proving Conjecture 1.13 is to establish analogous statements for heavy paths. Indeed, the method developed in [7] (and later used in [22] and [18]) for heavy paths is not applicable in the new setting.

Our main contribution in this paper is to develop a method to handle heavy paths for $t * S_{b,k}^s$ -free graphs, resulting in Lemma 3.1. We believe that some of the ideas we developed here can be further expanded to potentially yield further progress on Conjecture 1.1 and Conjecture 1.13.

3.2 Building $t * S_{b,k}^s$ using heavy paths

The rest of the section is devoted to proving Lemma 3.1. The proof consists of two parts: the case of $j > \frac{k+b}{2}$ (Lemma 3.3) and the case of $2 \leq j \leq \frac{k+b}{2}$ (Lemma 3.4). We would like to point out that it is possible to merge the two cases into one. However, we feel that each case contains useful ideas and prefer to keep them separate.

3.2.1 Long heavy paths: the $j > \frac{k+b}{2}$ case

Lemma 3.3. *Let G be a $t * S_{b,k}^s$ -free K -almost-regular graph on n vertices with minimum degree $\delta = \omega(1)$. Then provided that L is sufficiently large compared to s, t, k, K , for any $\frac{k+b}{2} < j \leq k$, the number of j -heavy paths in G is at most $\frac{n\delta^j}{L}$.*

Proof. We define some constants as follows. Let

$$c_1 = \frac{1}{4Lf(j-1, L)^2}, c_2 = \frac{c_1}{4K^b}, c_3 = \frac{c_1}{4K^{j-b}}, c_4 = \frac{c_3}{bK^{b-1}}, c_5 = \frac{c_2}{K^{j-b}}, c_6 = \left(\frac{c_5}{2}\right)^t \cdot c_2.$$

Suppose to the contrary that the number of j -heavy paths is at least $\frac{n\delta^j}{L}$. By averaging, there exists a vertex w such that the family \mathcal{P}_w consisting of all the j -heavy paths of the form $xx_1 \cdots x_{b-1}wx_{b+1} \cdots x_{j-1}y$ has size at least $\frac{\delta^j}{L}$. Let X be the set of vertices in G that play the role of x in some member of \mathcal{P}_w and Y the set of vertices in G that play the role of y in some member of \mathcal{P}_w . Then $X \subseteq \Gamma_b(w)$ and $Y \subseteq \Gamma_{j-b}(w)$. Since G is K -almost-regular and thus has maximum degree at most $K\delta$, we have

$$|X| \leq (K\delta)^b \text{ and } |Y| \leq (K\delta)^{j-b}. \tag{1}$$

Note that X, Y may not be disjoint. We define an auxiliary graph B on $X \cup Y$, such that $\forall x \in X, y \in Y, xy \in E(B)$ if and only if some member P of \mathcal{P}_w have ends x and y .

Claim 1. *For every $x \in X$ there is an (x, w) -path of length b in G . For every $y \in Y$ there is a (w, y) -path of length $j - b$ in G . For all $x \in X, y \in Y$ such that $xy \in E(B)$ there exist at least L internally disjoint (x, y) -paths of length j in G .*

Proof of Claim 1. The first two statements follow from the definitions of X and Y . Suppose $x \in X, y \in Y$ and $xy \in E(B)$. By definition, some member $P \in \mathcal{P}_w$ has x, y as ends. By the definition of \mathcal{P}_w , P is j -heavy and thus there exist at least $f(j, L)$ many j -admissible paths with ends x and y in G . By Lemma 2.6 among them we can find at least

$$f(j, L) / [j^2 f(j-1, L)^2] \geq L$$

that are pairwise vertex disjoint outside $\{x, y\}$, where the inequality holds by Proposition 2.2. □

For any fixed $x \in X$ and $y \in Y$, by Lemma 2.5 there are at most $f(b, L) \cdot f(j - b, L)$ members of \mathcal{P}_w that have ends x and y . Hence

$$e(B) \geq \frac{|\mathcal{P}_w|}{f(b, L)f(j-b, L)} \geq \frac{\delta^j}{Lf(j-1, L)^2}.$$

Now, let us colour each vertex in $X \cup Y$ with colour 1 or 2 independently at random with probability $\frac{1}{2}$ each. Let X_1 denote the set of vertices in X that receive colour 1 and Y_2 the set of vertices in Y that receive colour 2. Let \tilde{B} denote the subgraph of B consisting of edges that join a vertex in X_1 to a vertex in Y_2 . Each edge of B has probability at least $1/4$ of being in \tilde{B} . Hence there exists a colouring such that the resulting \tilde{B} has at least $(1/4)e(B)$ edges. Then \tilde{B} is bipartite

with parts X_1 and Y_2 and by our discussion

$$e(\tilde{B}) \geq \frac{\delta^j}{4Lf(j-1, L)^2} = c_1\delta^j. \tag{2}$$

By Lemma 2.10, \tilde{B} contains a subgraph B' with parts $X' \subseteq X_1$ and $Y' \subseteq Y_2$ such that

$$\forall x \in X', d_{B'}(x) \geq \frac{e(\tilde{B})}{4|X_1|} \geq \frac{c_1}{4K^b}\delta^{j-b} = c_2\delta^{j-b}, \tag{3}$$

and

$$\forall y \in Y', d_{B'}(y) \geq \frac{e(\tilde{B})}{4|Y_2|} \geq \frac{c_1}{4K^{j-b}}\delta^b = c_3\delta^b. \tag{4}$$

By (3) and (4),

$$|X'| \geq c_3\delta^b, \text{ and } |Y'| \geq c_2\delta^{j-b}.$$

Since $X' \subseteq X$, by Claim 1, there are at least $|X'|$ paths of length b with one end w and another end in X' . By Lemma 2.8, there exists a spider T of height b with centre w and leaves in X' whose number of legs is at least $\frac{|X'|}{b(K\delta)^{b-1}} \geq \frac{c_3}{bK^{b-1}}\delta = c_4\delta$.

Let $X'' \subseteq X'$ be the leaf set of this spider. Then $|X''| \geq c_4\delta$. Let B'' be the subgraph of B' induced by $X'' \cup Y'$. By (1) and (3)

$$e(B'') \geq c_2\delta^{j-b}|X''| \geq \frac{c_2}{K^{j-b}}|X''||Y'| = c_5|X''||Y'|.$$

Since $|X''| \geq c_4\delta$ and $\delta = \omega(1)$, for sufficiently large n we may assume that $c_5|X''| \geq 2t$. By Lemma 2.9

$$\exists X_0 \subseteq X'' \text{ such that } |X_0| = t \text{ and } |N_{B''}^*(X_0)| \geq (c_5/2)^t|Y'|. \tag{5}$$

Now, let us fix a t -set $X_0 \subseteq X''$ guaranteed in (5). Let T_0 be the sub-spider of T with leaf set X_0 . Let $Y'' = N_{B''}^*(X_0)$. Then

$$|Y''| \geq (c_5/2)^t|Y'| \geq (c_5/2)^t c_2\delta^{j-b} = c_6\delta^{j-b}.$$

Let \mathcal{C} be the family of paths of length $j - b$ with one end w and another end in Y'' . By Claim 1, $|\mathcal{C}| \geq |Y''| \geq c_6\delta^{j-b}$. Since G has maximum degree at most $K\delta$, for any vertex $u \neq w$, the number of paths in \mathcal{C} that contain u is at most $(j - b)(K\delta)^{j-b-1} < j(K\delta)^{j-b-1}$. Let \mathcal{C}_1 be the family of paths in \mathcal{C} that are vertex disjoint from $V(T_0) - \{w\}$. Then

$$|\mathcal{C}_1| \geq |\mathcal{C}| - (|V(T_0)| - 1)j(K\delta)^{j-b-1} \geq c_6\delta^{j-b} - btj(K\delta)^{j-b-1} \geq (c_6/2)\delta^{j-b},$$

where the last inequality holds for sufficiently large n because $\delta = \omega(1)$. As $j > \frac{k+b}{2}$, we have $k - j < j - b$. Applying Lemma 2.8 to \mathcal{C}_1 , as $\delta = \omega(1)$, there exists a t -legged spider T_1 of height $k - j$ with centre $v_1 \neq w$ and leaf set $Y_1 \subseteq Y''$. Note that $V(T_0) \cap V(T_1) = \emptyset$. Using the same strategy, we can find $s - 1$ vertex disjoint t -legged spiders T_1, \dots, T_{s-1} of height $k - j$ one by one, with T_i 's centre $v_i \neq w$ and leaf set $Y_i \subseteq Y''$, such that $V(T_i) \cap V(T_0) = \emptyset$.

Suppose $X_0 = \{x_1, \dots, x_t\}$. For each $i \in [s - 1]$, suppose $Y_i = \{y_1^i, \dots, y_t^i\}$. Let $Y_0 = \bigcup_{i=1}^{s-1} Y_i$. Then $Y_0 \subseteq Y'' = N_{B''}^*(X_0)$. Hence, $\forall x \in X_0, y \in Y_0, xy \in E(B'') \subseteq E(B)$ and by Claim 1 there exist at least L internally disjoint paths of length j joining x and y . As L is a sufficiently large constant, we can greedily find $t(s - 1)$ paths $P_{i,\ell}$ of length j , such that for any $i \in [t]$ and $\ell \in [s - 1]$, $P_{i,\ell}$ has ends x_i and y_ℓ^i and intersects $\bigcup_{i=0}^{s-1} V(T_i)$ only in x_i and y_ℓ^i and such that the $P_{i,\ell}$'s are pairwise vertex disjoint outside $\bigcup_{i=0}^{s-1} V(T_i)$. Now, $(\bigcup_{i=0}^{s-1} T_i) \cup (\bigcup_{i \in [t], \ell \in [s-1]} P_{i,\ell})$ forms a copy of $t * S_{b,k}^s$ in G , a contradiction. This completes our proof. \square

3.2.2 Short heavy paths: the $2 \leq j \leq \frac{k+b}{2}$ case

This subsection handles the most difficult part of our main proof and is where most of the new ideas are used.

Lemma 3.4. *Let G be a $t * S_{b,k}^s$ -free K -almost-regular graph on n vertices with minimum degree $\delta = \omega(1)$. Then provided that L is sufficiently large compared to s, t, k, K , for any $2 \leq j \leq \frac{k+b}{2}$, the number of j -heavy paths is at most $\frac{(j+1)^{j+1}}{L} n \delta^j$.*

We break the proof of Lemma 3.4 into several steps. The general strategy is to show that if the family \mathcal{F} of j -heavy paths is too large then we find a copy of $t * S_{b,k}^s$ in G , which is a contradiction. We start by doing some cleaning to \mathcal{F} in order to set up further arguments. Before that, let us set some constants to be used throughout the subsection.

Definition 3.5. Let $D = 2K^j L f(j-1, L)^2$ and $M = D^{s+1}$.

Comparing Definition 2.1 and Definition 3.5, we see that

$$f(j, L) = 10j^4 D^{s+3} = 10j^4 D^2 \cdot M. \tag{6}$$

Now we introduce our cleaning lemma. Given a path $P = v_0 v_1 \cdots v_j$ and $0 \leq i < j$, we define the *initial i -segment* of P to be the subpath $v_0 v_1 \cdots v_i$.

Lemma 3.6. *Let G be a K -almost-regular graph on n vertices with minimum degree $\delta = \omega(1)$. Suppose that the number of j -heavy paths is at least $\frac{(j+1)^{j+1}}{L} n \delta^j$. Then there exist a vertex w , vertex disjoint sets A_0, \dots, A_j and a family \mathcal{F} of j -heavy paths with $\bigcup_{i=0}^j A_i = \bigcup_{P \in \mathcal{F}} V(P)$ satisfying*

1. $A_0 \subseteq \Gamma_1(w)$ and $A_j \subseteq \Gamma_{j-1}(w)$.
2. Each member of \mathcal{F} has the form $v_0 v_1 \cdots v_j$ where $\forall i \in \{0, 1, \dots, j\}, v_i \in A_i$.
3. There exists a set V_0 with $A_0 \subseteq V_0 \subseteq \Gamma_1(w) \setminus A_j$ such that for every $y \in A_j$, there are at least $\frac{|V_0|}{D}$ many $x \in V_0$ such that x, y are ends of a heavy j -path in G . Furthermore, $|V_0| \geq (2K/D)\delta$.
4. For each $x \in A_0$, there are at least M vertices $y \in A_j$ such that x, y are ends of at least DM members of \mathcal{F} . For each $y \in A_j$, there are at least M vertices $x \in A_0$ such that x, y are ends of at least DM members of \mathcal{F} .
5. For each $P \in \mathcal{F}$ and $0 \leq i < j$, the initial i -segment of P is contained in at least $jM(K\delta)^{j-i-1}$ members of \mathcal{F} .

Proof. Let \mathcal{C} be the collection of all j -heavy paths in G . By our assumption, $|\mathcal{C}| \geq \frac{(j+1)^{j+1}}{L} n \delta^j$. Let us independently colour each vertex of G with a colour in $\{0, 1, \dots, j\}$, with each colour chosen uniformly at random. For each $0 \leq i \leq j$, let V'_i denote the set of vertices in G receiving colour i . For any j -heavy path $P = v_0 v_1 \cdots v_j$, call P *good* if $\forall 0 \leq i \leq j, v_i \in V'_i$. Let \mathcal{C}' denote the family of all good heavy j -paths. Clearly each j -heavy path in G is good with probability $(\frac{1}{j+1})^{j+1}$. So there exists a vertex colouring for which

$$|\mathcal{C}'| \geq \frac{|\mathcal{C}|}{(j+1)^{j+1}} \geq \frac{n \delta^j}{L}.$$

Let us fix such a colouring and the corresponding \mathcal{C}' .

By averaging, there exists a vertex w such that the subfamily \mathcal{P}_w of members of \mathcal{C}' of the form $v_0 w v_2 \cdots v_j$ has size at least $|\mathcal{P}_w| \geq \frac{\delta^j}{L}$. For each $i \in \{0, 1, \dots, j\}$, let V_i be the set of vertices in V'_i

that are contained in members of \mathcal{P}_w . By our definitions, $V_0 \subseteq \Gamma_1(w)$ and $V_j \subseteq \Gamma_{j-1}(w)$. Since G has maximum degree at most $K\delta$, we have

$$|V_0| \leq K\delta \text{ and } |V_j| \leq (K\delta)^{j-1}. \tag{7}$$

Let B denote the auxiliary bipartite graph with a bipartition (V_0, V_j) such that $\forall x \in V_0, y \in V_j, xy \in E(B)$ if and only if x, y are ends of some member of \mathcal{P}_w . For each $xy \in E(B)$ with $x \in V_0, y \in V_j$, let \mathcal{P}_{xy} be the subfamily of members of \mathcal{P}_w that cover x, y and let \mathcal{J}_{xy} be the family of j -heavy paths in G that have x, y as ends.

Claim 1. For each $xy \in E(B)$, we have $1 \leq |\mathcal{P}_{xy}| \leq f(j-1, L)$ and $|\mathcal{J}_{xy}| \geq f(j, L)$.

Proof of Claim 1. Let $xy \in E(B)$, with $x \in V_0, y \in V_j$. That $|\mathcal{P}_{xy}| \geq 1$ is clear. Let $P \in \mathcal{P}_{xy}$. By definition P is j -admissible and $P = xw \cup Q$, where Q is a (w, y) -path of length $j-1$. Since P is admissible, Q is $(j-1)$ -light. So the number of possible Q in G is at most $f(j-1, L)$. So, $|\mathcal{P}_{xy}| \leq f(j-1, L)$. Next, since P is a j -heavy path in G with ends x, y , by definition, G contains at least $f(j, L)$ j -heavy paths with ends x, y . So $|\mathcal{J}_{xy}| \geq f(j, L)$. \square

By Claim 1 and Definition 3.5

$$e(B) \geq \frac{|\mathcal{P}_w|}{f(j-1, L)} \geq \frac{\delta^j}{Lf(j-1, L)} \geq (2K^j/D)\delta^j. \tag{8}$$

Since $|V_j| \leq (K\delta)^{j-1}$, (8) implies

$$|V_0| \geq e(B)/|V_j| \geq (2K/D)\delta. \tag{9}$$

Let V_j^* be the set of $y \in V_j$ for which $d_B(y) \geq \frac{|V_0|}{D}$. Let B^* denote the subgraph of B induced by $V_0 \cup V_j^*$. Then

$$e(B^*) \geq e(B) - |V_0||V_j|/D \geq (2K^j/D)\delta^j - (K\delta)(K\delta)^{j-1}/D = K^j\delta^j/D. \tag{10}$$

For each $xy \in E(B^*)$, we have

$$|\mathcal{J}_{xy}| \geq f(j, L) \geq 10j^2D^2M,$$

where the last inequality holds by (6). Let \mathcal{J}'_{xy} be a subfamily of \mathcal{J}_{xy} of size exactly $10j^2D^2M$. Let

$$\mathcal{F}_0 = \bigcup_{xy \in E(B^*)} \mathcal{J}'_{xy}. \tag{11}$$

Then by (10)

$$|\mathcal{F}_0| = e(B^*) \cdot 10j^2D^2M \geq 10j^2DMK^j\delta^j. \tag{12}$$

We next obtain \mathcal{F} from \mathcal{F}_0 through some further cleaning. Initially let $\mathcal{F} = \mathcal{F}_0$. Throughout the process, for each $x \in V_0, y \in V_j^*$ let $\lambda(x, y)$ denote the number of remaining members of \mathcal{F} that have ends x, y . We update the function $\lambda(x, y)$ automatically after each removal. Whenever is a vertex $x \in V_0$ such that the number of $y \in V_j^*$ with $\lambda(x, y) \geq DM$ is less than M (which we refer to as x becomes small), remove all the members of \mathcal{F} that contain x . Similarly, whenever there is a vertex $y \in V_j^*$ such that the number of $x \in X$ with $\lambda(x, y) \geq DM$ is less than M (which we refer to as y becomes small), remove all the members of \mathcal{F} that contains y . Whenever there is a member $P \in \mathcal{F}$ (viewed as a path from V_0 to V_j^*) contains an initial i -segment I , for some $0 \leq i < j$, that is contained is less than $jM(K\delta)^{j-i-1}$ members of \mathcal{F} we remove all the members of \mathcal{F} containing I . We continue the process until no further removal can be performed.

The number of members of \mathcal{F} we removed for each $x \in V_0$ that becomes small is at most

$$\left(|V_j^*| - M\right) DM + M (10j^2D^2M) \leq 2DM(K\delta)^{j-1},$$

for sufficiently large n , since $\delta = \omega(1)$. Similarly, the number of members of \mathcal{F} that we removed for each vertex $y \in V_j^*$ that becomes small is at most

$$(|V_0| - M)DM + M(10j^2D^2M) \leq 2DM(K\delta).$$

So, the total number of members of \mathcal{F} we removed due to either a vertex in X becoming small or a vertex in Y^* becoming small is at most

$$|V_0| \cdot 2DM(K\delta)^{j-1} + |V_j^*| \cdot 2DM(K\delta) \leq 4DM(K\delta)^j.$$

The number of members of \mathcal{F} that we removed due to some initial segment is contained in too few members is at most

$$\sum_{i=0}^{j-1} |V_0|(K\delta)^i \cdot jM(K\delta)^{j-i-1} \leq j^2M(K\delta)^j.$$

Combining the above two inequalities, the total number of members of \mathcal{F} that we removed is at most

$$(j^2 + 4D)M(K\delta)^j \leq 5j^2DMK^j\delta^j \leq \frac{|\mathcal{F}_0|}{2}.$$

So in particular, the final \mathcal{F} is non-empty.

Now, for each $0 \leq i \leq j$, let A_i be the set of vertices in V_i that are contained in members of the final \mathcal{F} . In particular, note that $A_j \subseteq V_j^*$. Let us check that w, A_0, \dots, A_j and \mathcal{F} satisfy the five conditions of the lemma. Condition 1 and condition 2 clearly hold by our discussion so far. Condition 3 holds since $A_j \subseteq V_j^*$ and each vertex $y \in V_j^*$ satisfies $d_B(y) \geq \frac{|V_0|}{D}$ and $|V_0| \geq (2K/D)\delta$ by (9). Conditions 4 and 5 hold due to our cleaning rules. This completes the proof of the lemma. \square

Lemma 3.7. *Let G, A_0, \dots, A_j and \mathcal{F} be as stated in Lemma 3.6. Then*

1. *For any $0 \leq i < j$ and any $u \in A_i$, there exists an M -legged spider of height $j - i$ with centre u and leaves in A_j .*
2. *Let $F = \{uv : u \in A_{j-1}, v \in A_j \text{ and } \exists P \in \mathcal{F}, uv \in E(P)\}$. Then F has minimum degree at least M .*

Proof. Fix any i with $0 \leq i < j$ and $u \in A_i$. By the definition of A_i there exists $P = v_0v_1 \dots v_j \in \mathcal{F}$, where $v_i = u$. Let $I = v_0v_1 \dots v_i$. By condition 5 of Lemma 3.6, I is contained in at least $jM(K\delta)^{j-i-1}$ members of \mathcal{F} . In other words, the family $\mathcal{Q} = \{Q : Q \in A_i \times \dots \times A_j, I \cup Q \in \mathcal{F}\}$ has size at least $jM(K\delta)^{j-i-1}$. Since each member of \mathcal{Q} is a path of length $j - i$ from u to a vertex in A_j , by Lemma 2.8, there exists a spider of height $j - i$ with centre u and leaves in A_j whose number of legs is at least

$$\frac{jM(K\delta)^{j-i-1}}{j(K\delta)^{j-i-1}} = M.$$

This proves part 1 of the lemma. Applying part 1 with $i = j - 1$, we have $\forall u \in A_{j-1}, d_F(u) \geq M$. Let $y \in A_j$. By condition 4 of Lemma 3.6, there exist a vertex $x \in A_0$ such that there are at least DM members of \mathcal{F} that have ends x, y . By Lemma 2.6, among these there are at least $DM/[j^2f(j-1, L)^2] \geq M$ of them that are pairwise vertex disjoint outside $\{x, y\}$. In particular, this implies $d_F(y) \geq M$. So part 2 also holds. \square

Definition 3.8. For the rest of the subsection, we write $b = qj + b'$ where q and b' are integers with $1 \leq b' \leq j$.

The next lemma plays an important role in our proof of Lemma 3.4. It sets up a well-placed s -legged spider of height b to be used in building a copy of $t * S_{b,k}^s$.

Lemma 3.9. *Let G, w, A_0, \dots, A_j , and \mathcal{F} be as stated in Lemma 3.6. Let $N = M/D$.*

1. *If q is even, then there are N -legged spiders T and T' in G , both of height b such that the leaf set of T is contained in $A_{b'}$ and the leaf set of T' is contained in $A_{b'-1}$.*
2. *If q is odd, then there are N -legged spiders T and T' in G , both of height b , such that the leaf set of T is contained in $A_{j-b'}$ and the leaf set of T' is contained in $A_{j-b'+1}$.*

Proof. Let B be a bipartite graph with parts A_0 and A_j such that $\forall x \in A_0, y \in A_j \ xy \in E(B)$ if and only if at least DM members of \mathcal{F} have ends x, y . By Lemma 3.6 condition 4, B has minimum degree at least M . By Lemma 2.6, $\forall xy \in E(B)$, there exist at least $DM/[j^2f(j-1, L)^2] \geq M$ internally disjoint members of \mathcal{F} with ends x, y .

Fix a vertex $u \in A_0$. Since $\delta(B) \geq M = DN \geq (q+1)N$, where the last inequality follows from Definition 3.5, we can greedily grow an N -legged spider R in B that has centre u and height $q+1$. Since $\forall xy \in E(B)$ there are M internally disjoint members of \mathcal{F} with ends x, y , we can replace each edge ab of R with a member of \mathcal{F} with ends a, b so that the resulting graph is an N -legged spider S of height $(q+1)j$ in G . Let A be the set of vertices in S that are at distance $b = qj + b'$ from u in S . It is easy to see from the definition of S that if q is even then $A \subseteq A_{b'}$ and that if q is odd then $A \subseteq A_{j-b'}$. Let T be the sub-spider of S with centre u and leaf set A . Then T satisfies the first halves of statements 1 and 2.

Now, fix a subset $A'_0 \subseteq A_0$ of size N . Since B has minimum degree at least $M \geq (q+2)N + 1$, in B we can find N disjoint paths of length $q+1$, Q_1, \dots, Q_N , avoiding w , such that $\forall i \in [N]$, Q_i starts from a vertex $x_i \in A'_0$. By a similar reason as in the previous paragraph, we can replace the edges in $\bigcup_{i=1}^N Q_i$ by members of \mathcal{F} that avoid w such that for each $i \in [N]$, Q_i is turned into a path P_i of length $(q+1)j$ in G that still avoids w and that P_1, \dots, P_N are vertex disjoint. Let $S' = \bigcup_{i=1}^N P_i \cup \{wx_1, \dots, wx_N\}$. Then S' is an N -legged spider in G with centre w and height $(q+1)j+1$. Let A' be the set of vertices in S' that are distance $b = qj + b'$ from w in S' . It is easy to see by the definition of S' that if q is even then $A' \subseteq A_{b'-1}$ and that if q is odd then $A' \subseteq A_{j-b'+1}$. Let T' be the sub-spider of S' with centre w and leaf set A' . Then T' satisfies the second halves of statement 1 and 2. \square

Lemma 3.10. *Let G, w, A_0, \dots, A_j and \mathcal{F} be as stated in Lemma 3.6. Let m, m' be positive integers such that $m' \leq \frac{m}{2D}$ and $m \leq \frac{M}{2k}$. Let $0 \leq r \leq j$ and $U \subseteq A_r$ be a subset of size m . Let W be a vertex set such that $W \cap U = \emptyset$ and $|W| \leq M/2$. If $p := k + r - 2j$ is non-negative and even, then there exists an m' -legged spider T' with height k and leaf set $U' \subseteq U$ such that $V(T') \setminus U'$ is disjoint from $W \cup U$.*

Proof. Suppose $U = \{u_1, \dots, u_m\}$. Since $U \subseteq A_r$, by Lemma 3.7 statement 1, for each $i \in [m]$, there exists an M -legged spider of height $j-r$ with centre u_i and leaves in A_j . Since $M \geq km + |W| \geq (j-r+1)m + |W|$, by a greedy process, we can find a collection of vertex disjoint paths Q_1, \dots, Q_m , where for each $i \in [m]$, Q_i is a path of length $j-r$ joining u_i to a vertex y_i in A_j that avoids the set W . Let

$$Y = \{y_1, \dots, y_m\} \subseteq A_j.$$

By Lemma 3.7 statement 2, the graph

$$F = \{ab : a \in A_{j-1}, b \in A_j \text{ and } \exists P \in \mathcal{F}, ab \in E(P)\}$$

has minimum degree at least M . Using a greedy process we can find in F a collection of vertex disjoint paths R_1, \dots, R_m , where for each $i \in [m]$, R_i is a path in F of length p that joins y_i to some vertex z_i and avoids the set $(W \cup \bigcup_{\ell=1}^m V(Q_\ell)) \setminus \{y_i\}$. For each $i \in [m]$, since $y_i \in A_j$ and p is

even, $z_i \in A_j$ as well. Now $\{Q_i \cup R_i : i \in [m]\}$ is family of m disjoint paths of length $j - r + p$ that avoid W .

Let $Z = \{z_1, \dots, z_m\}$. Since $Z \subseteq A_j$, by Lemma 3.6 condition 3, there exists a set V_0 with $A_0 \subseteq V_0 \subseteq \Gamma_1(w) \setminus A_j$ such that $|V_0| \geq (2K/D)\delta$ and for each $i \in [m]$, there are at least $|V_0|/D$ many $x \in V_0$ such that x, z_i are the ends of a j -heavy path in G . Let B be a bipartite graph with parts V_0 and Z such that $\forall x \in V_0, z \in Z, xz \in E(B)$ if and only if x, z are the ends of a j -heavy path in G . Then $e(B) \geq |V_0||Z|/D$. Since $|Z|/D = m/D \geq 2m'$, by Lemma 2.9 there exists an m' -set $Z' \subseteq Z$ such that

$$|N_B^*(Z')| \geq \left(\frac{1}{2D}\right)^{m'} |V_0| \geq \left(\frac{1}{2D}\right)^M |V_0|.$$

Since $|V_0| \geq (K/2D)\delta$, while $\delta = \omega(1)$, for sufficiently large n , we have $|N_B^*(Z')| > |\bigcup_{i=1}^m V(Q_i \cup R_i) \cup W|$. So, there exists a vertex $x \in V_0 \setminus (\bigcup_{i=1}^m V(Q_i \cup R_i) \cup W)$ that is joined to all of Z' in B . Without loss of generality, suppose $Z' = \{z_1, \dots, z_{m'}\}$. For each $i \in [m']$, there exists a j -heavy path with ends x and z_i . In particular, by Lemma 2.6, there are at least $\frac{f(j,L)}{j^2 f(j-1,L)^2} \geq \frac{10j^4 D^2 M}{j^2 D} \geq M$ internally disjoint paths of length j between x and z_i , where the first inequality follows by Definition 3.5 and (6). As $M > m'k + |W|$, we can find paths $P_1, \dots, P_{m'}$, where $\forall i \in [m']$, P_i is a path of length j joining x to z_i such that $T' := \bigcup_{i=1}^{m'} (P_i \cup Q_i \cup R_i)$ is an m' -legged spider of height k with centre x and leaf set $\{u_1, \dots, u_{m'}\}$ and such that T' avoids $(U \setminus \{u_1, \dots, u_{m'}\}) \cup W$. The lemma holds for the above-defined T' and $U' = \{u_1, \dots, u_{m'}\}$. \square

Now, we are ready to prove Lemma 3.4.

Proof of Lemma 3.4. Suppose that the number of j -heavy paths in G is at least $\frac{(j+1)^{j+1}}{L} n \delta^j$. Let w, A_0, \dots, A_j and \mathcal{F} be obtained by Lemma 3.6. Our first step is to find an appropriate value of r to apply Lemma 3.10 to. Recall that $b = qj + b'$. Let

$$r := \begin{cases} b', & \text{if } q \text{ is even and } k + b' - 2j \text{ is even} \\ b' - 1, & \text{if } q \text{ is even and } k + b' - 2j \text{ is odd} \\ j - b', & \text{if } q \text{ is odd and } k + (j - b') - 2j \text{ is even} \\ j - b' + 1, & \text{if } q \text{ is odd and } k + (j - b') - 2j \text{ is odd} \end{cases}$$

As $1 \leq b' \leq j$, we have $0 \leq r \leq j$.

Let $p = k + r - 2j$. By the definitions of p and r , it is easy to see p is even. We claim that p is non-negative. To prove this, it is enough to show that $k + b' - 2j \geq 0$ when q is even, and that $k + (j - b') - 2j \geq 0$ when q is odd. First assume that q is even. If $q = 0$, then $b = b'$ and $k + b' - 2j = k + b - 2j \geq 0$ where the inequality holds by our assumption $j \leq \frac{k+b}{2}$; if $q \geq 2$, then $b \geq 2j + b'$ and thus $k + b' - 2j \geq b + b' - 2j \geq 2b' > 0$. Now assume q is odd. Then we have that $q \geq 1$ and thus $b \geq j + b'$. It follows that $k + (j - b') - 2j = k - (j + b') \geq b - (j + b') \geq 0$. Hence p is non-negative.

Now, let $m_1 = M/D$ and for $i = 2, \dots, s$, let $m_i = m_{i-1}/(2D)$. Using the definition of D , it is easy to check that $\forall i \in [s], m_i \leq M/(2sk)$. By Lemma 3.9, there exists an m_1 -legged spider T_1 with height b and leaf set $U_1 \subseteq A_r$. The idea of the rest of the proof is to apply Lemma 3.10 $s - 1$ times. Initially let $W = V(T_1) \setminus U_1$. Since $p = k + r - 2j$ is non-negative and even, applying Lemma 3.10 with m_1 and m_2 playing the roles of m and m' respectively and U_1 playing the role of U , we can find an m_2 -legged spider T_2 with height k and leaf set $U_2 \subseteq U_1$ such that $V(T_2) \setminus U_2$ is disjoint from $W \cup U_1$. Now, we add $V(T_2) \setminus U_2$ to W . Next, applying Lemma 3.10 with m_2, m_3 playing the roles of m and m' respectively and U_2 playing the role of U , we can find an m_3 -legged spider T_3 with height k and leaf set $U_3 \subseteq U_2$, such that $V(T_3) \setminus U_3$ is disjoint from $W \cup U_2$. Now, we add

$V(T_3) \setminus U_3$ to W . We continue like this. It is easy to check that we can carry out the process for at least $s - 1$ steps to find T_2, \dots, T_s . Indeed, within the first $s - 1$ steps W has size at most $km_1 + km_2 + \dots + km_{s-1} < 2km_1 = 2kM/D < M/2$. This together with the definitions of m_1, \dots, m_s ensures that the conditions of Lemma 3.10 are satisfied. But now $\bigcup_{i=1}^s T_i$ forms a copy of $t * S_{b,k}^s$ in G , a contradiction. This completes our proof of Lemma 3.4. \square

3.2.3 Proof of Lemma 3.1

Now we are in a position to prove Lemma 3.1.

Proof of Lemma 3.1. By Lemmas 3.3 and 3.4, for any $2 \leq j \leq k$, the number of j -heavy paths in G is at most $\max \left\{ \frac{n\delta^j}{L}, \frac{(j+1)^{j+1}}{L} n\delta^j \right\} = \frac{(j+1)^{j+1}}{L} n\delta^j$. This completes the proof. \square

4. Concluding remarks

In [25], Kang, Kim and Liu extended the definition of balanced rooted trees to that of balanced rooted bipartite graphs as follows. Let F be a bipartite graph and R a proper subset of $V(F)$ called the set of roots. For each non-empty set $S \subseteq V(F)$, let $\rho_F(S) = \frac{e_S}{|S|}$, where e_S is the number of edges in G with at least one end in S . Let $\rho(F) = \rho_F(V(F) \setminus R)$. We say that (F, R) is balanced if $\rho_F(S) \geq \rho(F)$ for every non-empty subset $S \subseteq V(F) \setminus R$. A real number $r \in (1, 2)$ is called *balancedly realizable* if there is a connected bipartite graph F and a set $R \subseteq V(F)$ such that (F, R) is balanced with $\rho_F = \frac{1}{2-r}$ and that there is a positive integer t_0 such that for all integers $t \geq t_0$, $\text{ex}(n, t * F) = \Theta(n^r)$ holds. By definition, a balancedly realizable number is a Turán exponent. Using a result of Erdős and Simonovits [11], Kang, Kim and Liu [25] proved the following.

Lemma 4.1 ([25]). *Let $a < b$ be two integers. If $2 - \frac{a}{b}$ is balancedly realizable, then $2 - \frac{a}{a+b}$ is also balancedly realizable.*

Proof of Corollary 1.4 and Corollary 1.5. By Theorem 1.9 and Theorem 1.15, for any positive integers p, k, b with $k \geq b$, $1 + \frac{p}{kp+b}$ is balancedly realizable. Corollary 1.4 follows by applying Lemma 4.1 repeatedly.

Now, suppose that $q = sp + p'$ where s is a positive integer and $0 \leq p' \leq \sqrt{p}$. Since it is known that $2 - \frac{1}{s}$ is a Turán exponent for all any integer $s \geq 2$, we may assume $p' > 0$. Now, as $0 < p' \leq \sqrt{p}$, there exists integers k and b such that $p = kp' + b$ and $k \geq p' - 1$ and $p' \geq b \geq 1$. Then $2 - \frac{p}{q} = 2 - \frac{kp'+b}{s(kp'+b)+p'}$. By Corollary 1.4, it follows that $2 - \frac{p}{q}$ is a Turán exponent. \square

Finally, even though we obtained all the Turán exponents that Janzer’s conjecture would give, it would still be very interesting to resolve his conjecture in the full. For the general bipartite Turán problem, while tools such as dependent random choice have found success in the denser end of the spectrum for bipartite graphs, it would be very interesting to develop more tools for the sparser end of the spectrum. The recent active study of the Turán problem for subdivisions is a step in that direction. It will be very interesting to continue explore problems of such nature.

References

- [1] Alon, N., Krivelevich, M. and Sudakov, B. (2003) Turán numbers of bipartite graphs and related Ramsey-type questions. *Combin. Probab. Comput.* **12**(5-6) 477–494.
- [2] Bukh, B. (2015) Random algebraic construction of extremal graphs. *Bull. Lond. Math. Soc.* **47**(6) 939–945.
- [3] Bukh, B. and Conlon, D. (2018) Rational exponents in extremal graph theory. *J. Eur. Math. Soc.* **20** 1747–1757.
- [4] Brown, W. G. (1966) On graphs that do not contain a Thomsen graph. *Can. Math. Bull.* **9**(3) 281–285.
- [5] Conlon, D. (2019) Graphs with few paths of prescribed length between any two vertices. *Bull. Lond. Math. Soc.* **51**(6) 1015–1021.

- [6] Conlon, D. and Lee, J. (2021) On the extremal number of subdivisions. *Int. Math. Res. Not.* **2021**(12) 9122–9145.
- [7] Conlon, D., Janzer, O. and Lee, J. (2021) More on the extremal number of subdivisions. *Combinatorica* **41** 465–494.
- [8] Erdős, P. (1986) Problems and results in combinatorial analysis and graph theory. *Proc. First Jpn. Conf. Graph Theory Appl. (Hakone)* **72**(1988) 81–92.
- [9] Erdős P. and Stone A. H. (1946) On the structure of linear graphs. *Bull. Am. Math. Soc* **52**(12) 1087–1091.
- [10] Erdős, P. and Simonovits, M. (1966) A limit theorem in graph theory. *Studia Sci. Math. Hungar.* **1** 51–57.
- [11] Erdős, P. and Simonovits, M. (1970) Some extremal problems in graph theory. *Comb. Theory Appl. (Proc. Colloq. Balatonfüred 1969)* **1** 370–390.
- [12] Erdős, P. and Stone, H. (1946) On the structure of linear graphs. *Bull. Am. Math. Soc.* **52** 1087–1091.
- [13] Faudree, R. and Simonovits, M. (1983) On a class of degenerate extremal graph problems. *Combinatorica* **3** 83–93.
- [14] Frankl, P. (1986) All rationals occur as exponents. *J. Combin. Theory Ser. A* **42** 200–206.
- [15] Füredi, Z. (1991) On a Turán type problem of Erdős. *Combinatorica* **11** 75–79.
- [16] Füredi, Z. and Simonovits, M. (2013) The history of the degenerate (bipartite) extremal graph problems. *Erdős centennial, Bolyai Soc. Math. Stud.* **25** 169–264, János Bolyai Math. Soc., Budapest, 2013. See also arXiv:1306.5167.
- [17] Janzer, O. (2019) Improved bounds for the extremal number of subdivisions. *Electron. J. Comb.* **26**(3) P3.3.
- [18] Janzer, O. (2020) The extremal number of the subdivisions of the complete bipartite graph. *SIAM J. Discrete Math.* **34**(1) 241–250.
- [19] Janzer, O. (2021) The extremal number of longer subdivisions. *Bull. London Math. Soc.* **53**(1) 108–118.
- [20] Jiang, T. (2011) Compact topological minors in graphs. *J. Graph Theory* **67**(2) 139–152.
- [21] Jiang, T., Ma, J. and Yepremyan, L. (2022) On Turán exponents of bipartite graphs. *Combin. Probab. Comput.* **31**(2) 333–344.
- [22] Jiang, T. and Qiu, Y. (2020) Turán numbers of bipartite subdivisions. *SIAM J. Discrete Math.* **34**(1) 556–570.
- [23] Jiang, T. and Qiu, Y. () Many Turán exponents via subdivisions, arXiv:1908.02385v1,
- [24] Jiang, T. and Seiver, R. (2012) Turán numbers of subdivided graphs. *SIAM J. Discrete Math* **26**(3) 1238–1255.
- [25] Kang, D. Y., Kim, J. and Liu, H. (2021) On the rational Turán exponent conjecture. *J. Combin. Theory Ser. B* **148** 149–172.
- [26] Kővári, T., Sós, V. T. and Turán, P. (1954) On a problem of K Zarankiewicz. *Colloq. Math.* **3**(1) 50–57.